arXiv:math/9912047v1 [math.CO] 6 Dec 1999

Combinatorial Properties of the Family of Maximum Stable Sets of a Graph

Vadim E. Levit and Eugen Mandrescu
Department of Computer Science
Holon Academic Institute of Technology
52 Golomb Str., P.O. Box 305
Holon 58102, ISRAEL
{levitv, eugen_m}@barley.cteh.ac.il

Abstract

The *stability number* $\alpha(G)$ of a graph G is the size of a *maximum stable set* of G, $core(G) = \cap\{S : S\ is\ a\ maximum\ stable\ in\ G\}$, and $\xi(G) = |core(G)|$. In this paper we prove that for a graph G without isolated vertices, the following assertions are true: (i) if $\xi(G) \leq 1$, then G is quasi-regularizable; (ii) if G is of order n and $\alpha(G) > (n + k - 1)/2$, for some $k \geq 1$, then $\xi(G) \geq k + 1$, and $\xi(G) \geq k + 2$, whenever $n + k - 1$ is even. The last finding is a strengthening of a result of Hammer, Hansen, and Simeone, which states that $\alpha(G) > n/2$ implies $\xi(G) \geq 1$. In the case of König-Egerváry graphs, i.e., for graphs enjoying $\alpha(G) + \mu(G) = n$, where $\mu(G)$ is the maximum size of a matching of G, we prove that $|core(G)| > |N(core(G))|$ is a necessary and sufficient condition for $\alpha(G) > n/2$. Moreover, for bipartite graphs without isolated vertices, $\xi(G) \geq 2$ is equivalent to $\alpha(G) > n/2$. We also show that Hall's marriage Theorem is valid for König-Egerváry graphs, and, it is sufficient to check Hall's condition only for one specific stable set, namely, for $core(G)$.

1 Introduction

Throughout this paper $G = (V, E)$ is a simple (i.e., a finite, undirected, loopless and without multiple edges) graph with vertex set $V = V(G)$ and edge set $E = E(G)$. If $X \subset V$, then $G[X]$ is the subgraph of G spanned by X. By $G - W$ we mean either the subgraph $G[V - W]$, if $W \subset V(G)$, or the partial subgraph $H = (V, E - W)$ of G, for $W \subset E(G)$. Anyway, we use $G - w$, whenever $W = \{w\}$. If $A, B \subset V$ and $A \cap B = \emptyset$, then (A, B) stands for the set $\{e = ab : a \in A, b \in B, e \in E\}$. A stable set S of maximum size will be referred as to a *maximum stable set* of G, and $\alpha(G) = |S|$ is the *stability number* of G. Let $\Omega(G)$ and $core(G)$ denote respectively the sets $\{S : S\ is\ a\ maximum\ stable\ set\ of\ G\}$ and $\cap\{S : S \in \Omega(G)\}$, while $\xi(G) = |core(G)|$. Clearly, any isolated vertex of a graph G is contained in $core(G)$. Let us define $isol(G)$ as the set of isolated vertices of G. The neighborhood of a vertex $v \in V$ is the set

$N(v) = \{w : w \in V \ and \ vw \in E\}$, and for $A \subset V, N(A) = \cup\{N(v) : v \in A\}$, while $N[A] = A \cup N(A)$. By C_n, K_n, P_n we denote the chordless cycle on $n \geq 4$ vertices, the complete graph on $n \geq 1$ vertices, and respectively the chordless path on $n \geq 3$ vertices.

A *matching* is a set of non-incident edges of G; a matching of maximum cardinality $\mu(G)$ is a *maximum matching*, and a *perfect matching* is a matching covering all the vertices of G. G is a *König-Egerváry graph* provided $\alpha(G) + \mu(G) = |V(G)|$, [2], [13]. According to a well-known result of König, [7], and Egerváry, [3], any bipartite graph enjoys this property.

A graph G is *α^+-stable* if $\alpha(G+e) = \alpha(G)$, for any edge $e \in E(\overline{G})$, where $\overline{G}$ is the complement of G, [4]. The following characterization of α^+-stable graphs, without any referring to this notion, had been proved in [5] three years before the above definition was proposed.

Theorem 1.1 *[5] A graph G is α^+-stable if and only if $\xi(G) \leq 1$.*

This result motivates that a graph G is referred to as: (i) α_0^+-stable, if $\xi(G) = 0$, and (ii) α_1^+-stable provided $\xi(G) = 1$, [11]. For instance, C_4 is α_0^+-stable, $K_3 + e$ is α_1^+-stable, and the diamond, i.e., the graph $K_4 - e$, is not α^+-stable (see Figure 1).


(a) (b)

Figure 1: Two non-α_0^+-stable graphs: (a) $K_4 - e$; (b) $K_3 + e$.

A graph G is *quasi-regularizable* if one can replace each edge of G with a non-negative integer number of parallel copies, so as to obtain a regular multigraph of degree $\neq 0$, [1]. For instance, the diamond is quasi-regularizable, while P_3 is not quasi-regularizable. Clearly, any quasi-regularizable graph has no isolated vertices. Moreover, a disconnected graph is quasi-regularizable if and only if any of its connected components is a quasi-regularizable graph.

In this paper we analyze the relationship between $\alpha(G)$ and $\xi(G)$. We show that if G has $|isol(G)| \neq 1$ and $\alpha(G) > (|V(G)|+k-1)/2$, then necessarily $\xi(G) \geq k+1$ holds; moreover, $\xi(G) \geq k+2$ is valid, whenever $|V(G)|+k-1$ is an even number. For $k = 1$, we obtain a strengthening of a result of Hammer, Hansen and Simeone, [6], which claims that $\xi(G) \geq 1$, whenever $\alpha(G) > |V(G)|/2$. The fact that $\alpha(G) > |V(G)|/2$ together with $|isol(G)| \neq 1$ implies $\xi(G) \geq 2$ were first established in [9] and [10] for bipartite graphs and König-Egerváry graphs respectively. From the historical perspective it is also worth mentioning that $\xi(T) \neq 1$ holds for any tree T of order at least two, as Gunther et al., [4], and independently, Zito, [14], have shown. Moreover, $\xi(G) \neq 1$ is true for an arbitrary bipartite graph G, [9].

We also thoroughly investigate the special cases of König-Egerváry graphs and bipartite graphs, for which some sufficient conditions implying $\alpha(G) > |V(G)|/2$ are found. We infer that Hall's marriage Theorem is true for König-Egerváry graphs as well as for bipartite graphs, and obtain a new characterization of König-Egerváry graphs having a perfect matching in terms of properties of $core(G)$.

2 The main result: $\alpha(G) > (n+k-1)/2 \Rightarrow \xi(G) \geq k+1$

Recall the following characterization of quasi-regularizable graphs, due to Berge.

Theorem 2.1 *[1] A graph G is quasi-regularizable if and only if $|S| \leq |N(S)|$ holds for any stable set S of G.*

Corollary 2.2 *If G is a quasi-regularizable graph, then $\alpha(G) \leq |V(G)|/2$.*

Lemma 2.3 *For a graph G let S_0 be a set of vertices such that*

$$|S_0| = \min\{|S| : S \text{ is stable in } G \text{ and } |S| > |N(S)|\}.$$

If $isol(G) = \emptyset$, then the following assertions hold:
(i) G is quasi-regularizable if and only if $|S_0| = 0$;
(ii) G is not quasi-regularizable if and only if $|S_0| \geq 2$.

Proof. (i) It is clear, according to Theorem 2.1. Consequently, G is not quasi-regularizable if and only if $|S_0| \geq 1$, in fact, if and only if $|S_0| \geq 2$, since S_0 is contained in some connected component of G. ■



Figure 2: Non-Quasi-Regularizable graphs.

If $isol(G) \neq \emptyset$, then the above set S_0 has cardinality $|S_0| = 1$. It is worth mentioning that there is a close relation between S_0 and $core(G)$. In order to see this we need the following characterization of a maximum stable set of a graph, due to Berge.

Theorem 2.4 *[1] A stable set S belongs to $\Omega(G)$ if and only if every stable set A of G, disjoint from S, can be matched into S.*

Lemma 2.5 *If G is not quasi-regularizable and S_0 is a set of vertices such that*

$$|S_0| = \min\{|S| : S \text{ is stable in } G \text{ and } |S| > |N(S)|\},$$

then $S_0 \subseteq core(G)$.

Proof. Clearly, the assertion is true for the case that $isol(G) \neq \emptyset$, since all the isolated vertices of G are contained in any $S \in \Omega(G)$, and S_0 consists of a single isolated vertex.

Now, let G be without isolated vertices, and suppose, on the contrary, that there exists some $S \in \Omega(G)$, which does not include S_0.

Case 1. $S_0 \cap S = \emptyset$. According to Theorem 2.4, S_0 can be matched into S, and this yields $|S_0| \leq |N(S_0)|$, in contradiction with the definition of S_0.

Case 2. $S_0 \cap S = S_1 \neq \emptyset$. Since $S_0 - S_1$ is stable and disjoint from $S \in \Omega(G)$, Theorem 2.4 implies that $S_0 - S_1$ can be matched into S. Hence, $|S_0 - S_1| \leq |N(S_0 - S_1)|$. Consequently, by $|S_0| > |N(S_0)|$ we get that $|S_1| > |N(S_1)|$, thus contradicting the minimality of S_0.

Thus, we may conclude that $S_0 \subseteq S$, for any $S \in \Omega(G)$, and this ensures that $S_0 \subseteq core(G)$. ■

Remark 1 *If G is not quasi-regularizable and satisfies $isol(G) = \emptyset$, then the inclusion in Lemma 2.5 may be sometimes strict. For instance, the graph G_1 in Figure 2 has $S_0 = core(G_1) = \{a, b\}$, while for the graph G_2 in the same Figure we note that $S_0 = \{a, b\} \subset core(G_2) = \{a, b, c\}$.*

Proposition 2.6 *Any α^+-stable graph free of isolated vertices is quasi-regularizable.*

Proof. Assume that G is a non-quasi-regularizable α^+-stable graph with $isol(G) = \emptyset$. According to lemmas 2.3 and 2.5, there exists some stable set S_0 in G, with $|S_0| \geq 2$, such that $S_0 \subseteq core(G)$. Consequently, we obtain that $\xi(G) \geq 2$, a contradiction, since by virtue of Theorem 1.1, G must satisfy the condition $\xi(G) \leq 1$. ■

The restriction "*free of isolated vertices*" in the proposition above is essential, since no graph G with $isol(G) \neq \emptyset$ can be quasi-regularizable, but there exist α^+-stable graphs having isolated vertices; e.g., any graph consisting of one isolated vertex and a $K_n, n \geq 2$, is α^+-stable. Nevertheless, $|isol(G)| \leq 1$ holds for any α^+-stable graph G.

Combining Proposition 2.6 and Corollary 2.2 we obtain the following:

Corollary 2.7 *If G is an α^+-stable graph with $isol(G) = \emptyset$, then $\alpha(G) \leq |V(G)|/2$.*

Remark 2 *The inequality in Corollary 2.7 is not true for any α^+-stable graph. For instance, the graph G consisting of two connected components, namely one isolated vertex and a C_4, is α^+-stable and $\alpha(G) = 3 > |V(G)|/2 = 5 / 2$. If G consists of one isolated vertex and a K_n, $n \geq 2$, then G is α^+-stable and $\alpha(G) = 2 > |V(G)|/2$, for $n = 2$, $\alpha(G) = 2 = |V(G)|/2$, for $n = 3$, while for $n \geq 4$, $\alpha(G) = 2 < |V(G)|/2$.*

Proposition 2.8 *Let G be a graph with $\alpha(G) > |V(G)|/2$. Then the following statements are equivalent:*

(i) G is α^+-stable;

(ii) G is α_1^+-stable;

(iii) G has a unique isolated vertex v, $\xi(G-v) = 0$ and $\alpha(G-v) = (|V(G)|-1)/2$.

Proof. If $|V(G)| = 1$, the result is obvious. Suppose now that $n = |V(G)| > 1$.
$(i) \Rightarrow (ii)$, (iii) As an α^+-stable graph, G may have at most one isolated vertex, and using now Corollary 2.7, we get that G has exactly one isolated vertex, say v. It follows that G is α_1^+-stable, $core(G) = \{v\}$ and $G - v$ is α_0^+-stable with $\alpha(G-v) = \alpha(G) - 1$. Since $G - v$ is α^+-stable and has no isolated vertices, Corollary 2.7 ensures that $\alpha(G-v) \leq (n-1)/2$. Hence, we obtain:

$$n/2 - 1 < \alpha(G) - 1 = \alpha(G-v) \leq (n-1)/2,$$

which implies that n must be odd and $\alpha(G-v) = (n-1)/2$.

The implications $(iii) \Rightarrow (ii)$ and $(ii) \Rightarrow (i)$ are clear. ■

Corollary 2.9 *There is no graph G satisfying $\alpha(G) > |V(G)|/2$, such that either (i) G is α_1^+-stable and of even order, or (ii) G is α_0^+-stable.*

Corollary 2.10 *If $\alpha(G) > |V(G)|/2$ and $\xi(G) = 1$, then $|V(G)| \equiv 1$ mod 2.*

Theorem 2.11 *Let G be a graph with $\alpha(G) > |V(G)|/2$. Then the following statements are true:*
(i) if $|isol(G)| \neq 1$, then $\xi(G) \geq 2$;
(ii) if $\xi(G) = 1$, then $|isol(G)| = 1$;
(iii) $\xi(G) = 0$ never holds.

Proof. (i) The result is clear, whenever $|isol(G)| \geq 2$, because $\xi(G) \geq |isol(G)|$. Let $isol(G) = \emptyset$, and suppose, on the contrary, that $\xi(G) \leq 1$. By Theorem 1.1, we infer that G is an α^+-stable graph. Hence, Corollary 2.7 ensures that $\alpha(G) \leq |V(G)|/2$, in contradiction with the premise on G. Therefore, $\xi(G) \geq 2$ is true.

(ii) If $\xi(G) = 1$, then $|isol(G)| = 1$, since otherwise, according to (i), G must satisfy $\xi(G) \geq 2$.

(iii) If $\xi(G) = 0$, then G is α^+-stable. Since $\alpha(G) > |V(G)|/2$, Proposition 2.8 implies that G must be α_1^+-stable, i.e., $\xi(G) = 1$, in contradiction with the assumption on G. ■

Remark 3 *The condition $\xi(G) \geq 2$ is not strong enough to ensure $\alpha(G) > n/2$. Moreover, for any positive integer k, one can choose an arbitrarily large positive integer p and build a graph $G = (V, E)$, with $n = k + p$ vertices, such that $\xi(G) = k$ and $\alpha(G) < n/2$. For instance, such a graph $G = (V, E)$ is defined by:*

$$V = \{x_i : 1 \leq i \leq k\} \cup \{y_i : 1 \leq i \leq p\},\ E = \{x_i y_1 : 1 \leq i \leq k\} \cup \{y_i y_j : 1 \leq i < j \leq p\}.$$

Theorem 2.11 is a strengthening of the following result, due to Hammer, Hansen and Simeone:

Corollary 2.12 *[6] If G has $\alpha(G) > |V(G)|/2$, then $\xi(G) \geq 1$.*

Proposition 2.13 *If* $G = (V, E)$ *and* $H = G[V - N[core(G)]]$, *then the following assertions are true:*

(i) H *has no isolated vertices;*

(ii) $\alpha(H) = \alpha(G) - \xi(G)$*;*

(iii) $S_H \in \Omega(H)$ *if and only if there is* $S_G \in \Omega(G)$*, such that* $S_H = S_G \cap V(H)$*;*

(iv) H *is* α_0^+*-stable;*

(v) $|S - core(G)| \leq |N(S) - N(core(G))|$ *holds for any* $S \in \Omega(G)$.

Proof. If $core(G) = \emptyset$, then $H = G$ and all the assertions are valid. Assume that $core(G) \neq \emptyset$. Let $S \in \Omega(G), A = S - core(G)$ and $B = V - S - N(core(G))$.

(i) Suppose, on the contrary, that H has an isolated vertex v.

Case 1. $v \in B$. Then $(\{v\}, core(G)) = (\{v\}, A) = \emptyset$, because $v \notin N(core(G))$ and it is isolated in H. Hence, $S \cup \{v\}$ is stable in G, contradicting the maximality of S.

Case 2. $v \in A$. Let $W \in \Omega(G), W \neq S$. Then $core(G) \subset W, W \cap B \neq \emptyset, W \cap N(core(G)) = \emptyset$ and maybe $W \cap A \neq \emptyset$. Hence, v is adjacent to no vertex in W, and since W is a maximum stable set in G, we infer that $v \in W$. It follows that $v \in core(G)$, in contradiction with the fact that $v \in A = S - core(G)$.

(ii) Since A is stable, it follows that $\alpha(G) - \xi(G) = |A| \leq \alpha(H)$. Suppose, on the contrary, that $\alpha(H) = |S_H| > |A|$, with $S_H \in \Omega(H)$. Hence, we get that: $(S_H \cap A, core(G)) = (S_H \cap B, core(G)) = \emptyset$, because $S_H \cap A \subset A \subset S$ and also $B \cap N(core(G)) = \emptyset$. Consequently, $S_H \cup core(G)$ is stable in G and $|S_H \cup core(G)| = |S_H| + \xi(G) > |A| + \xi(G) = \alpha(G)$, in contradiction with definition of $\alpha(G)$.

(iii) If $S_H \in \Omega(H)$, then $(S_H \cap B, core(G)) = \emptyset$, because $B \cap N(core(G)) = \emptyset$, and evidently $(S_H \cap A, core(G)) = \emptyset$. Hence, $S_G = core(G) \cup S_H$ is stable in G and since, by (ii), $\alpha(H) = |S_H| = \alpha(G) - \xi(G)$, we infer that $S_G \in \Omega(G)$.

Conversely, if $S_H = S_G \cap V(H)$ and $S_G \in \Omega(G)$, then S_H is stable in H as well, and $|S_H| = |S_G| - \xi(G) = \alpha(G) - \xi(G) = \alpha(H)$. Therefore, $S_H \in \Omega(H)$.

(iv) According to (iii), it follows that $core(H) = \emptyset$, i.e., H is α_0^+-stable.

(v) H is α_0^+-stable and has no isolated vertices. Hence, Corollary 2.7 implies that $\alpha(H) \leq |V(H)|/2$, i.e., $|S - core(G)| = |A| \leq |B| = |N(S) - N(core(G))|$. ■

Remark 4 *The inequality* $|S - A| \leq |N(S) - N(A)|$ *is not generally true for any subset* A *of a maximum stable set* S*. For instance, for the graph in Figure 3, one can take* $S = \{a, b, c, d, e\}, A = \{b, c, d\}$*, and get* $|S - A| = 2 > |N(S) - N(A)| = 1$.



Figure 3: A counterexample to $|S - A| \leq |N(S) - N(A)|$.

By Corollary 2.2, if $\alpha(G) > |V(G)|/2$, then G is not quasi-regularizable. From the point of view of Berge's Theorem 2.1, an elementary obstacle to being quasi-regularizable is the presence of a stable set S in G satisfying $|S| > |N(S)|$. An interesting question is how to present at least one such stable set for the graph G. The following result gives a canonical procedure for constructing such a stable set, when $\alpha(G) > |V(G)|/2$.

Theorem 2.14 *If a graph G has $\alpha(G) > |V(G)|/2$, then $|core(G)| > |N(core(G))|$.*

Proof. Let $S \in \Omega(G)$, $A = S - core(G), B = V - S - N(core(G))$, and also $H = G[V - N[core(G)])]$. By Proposition 2.13, $\alpha(H) = \alpha(G) - \xi(G) = |A|$. Suppose, on the contrary, that $\xi(G) \leq |N(core(G))|$. Since $\alpha(G) > |V(G)|/2$ we get $\alpha(G) = \xi(G) + |A| > |N(core(G))| + |B|$, which together with $\xi(G) \leq |N(core(G))|$ gives $|A| = \alpha(G) - \xi(G) > |V(G)|/2 - |N(core(G))| = |B|$. Therefore, it follows $\alpha(H) = |A| > |V(H)|/2$. Using again Proposition 2.13, we infer that H is α_0^+-stable and $isol(H) = \emptyset$. According to Corollary 2.7, it follows that $\alpha(H) \leq |V(H)|/2$, which contradicts the premise of the theorem. ■

The converse of Theorem 2.14 is not generally true. The graph in Figure 4 illustrates this assertion.



Figure 4: $\xi(G) > |N(core(G))|$ and $\alpha(G) < |V(G)|/2$.

Proposition 2.15 *If a graph G has $\alpha(G) > |V(G)|/2$, and $\xi(G) \leq k$, for some $k \geq 2$, then $\alpha(G) \leq (|V(G)| + k - 1)/2$.*

Proof. The result is clear for $|V(G)| = 1$.
If $|V(G)| > 1$, let us denote $H = G[V(G) - N[core(G)]]$. Then we obtain:

$$|V(H)| = |V(G)| - |core(G)| - |N(core(G))| = |V(G)| - \xi(G) - |N(core(G))|.$$

By Proposition 2.13, H is α_0^+-stable and $isol(H) = \emptyset$. According to Corollary 2.7, it follows that $\alpha(H) \leq |V(H)|/2 \leq (|V(G)| - \xi(G) - 1)/2$, and consequently, we infer that: $\alpha(G) = \alpha(H) + \xi(G) \leq (|V(G)| - \xi(G) - 1)/2 + \xi(G) = (|V(G)| + \xi(G) - 1)/2 \leq (|V(G)| + k - 1)/2$. ■

Theorem 2.16 *If $|isol(G)| \neq 1$ and $\alpha(G) > (|V(G)| + k - 1)/2$, for some $k \geq 1$, then the following assertions are true:*
(i) $\xi(G) \geq k + 1$, whenever $|V(G)| + k - 1$ is odd;
(ii) $\xi(G) \geq k + 2$, whenever $|V(G)| + k - 1$ is even.

Proof. Clearly, $n = |V(G)| \neq 1$. First we show that (i) is always valid. For $k = 1$ this assertion is exactly the claim of Theorem 2.11(i). Suppose that $k \geq 2$, and $\alpha(G) > (n + k - 1)/2 \geq n/2$. If $\xi(G) \leq k$, then by Proposition 2.15, it follows that $\alpha(G) \leq (n + k - 1)/2$, in contradiction with the premises on $\alpha(G)$. Therefore, $\xi(G) \geq k + 1$ is true for any $k \geq 1$. Assume that (ii) is not valid. Therefore we get that $\xi(G) \leq k + 1$, and because $\alpha(G) > (n + k - 1)/2 \geq n/2$, Proposition 2.15 implies that $\alpha(G) \leq (n + k + 1 - 1)/2 = (n + k)/2$. Hence we obtain the following contradiction:

$$(n+k-1)/2 < \alpha(G) \leq (n+k)/2,$$

since $\alpha(G)$ must be a positive integer. ■

For $k=1$, we obtain the following strengthening of Corollary 2.12 due to Hammer, Hansen and Simeone, and of Theorem 2.11(i):

Corollary 2.17 *If* $|isol(G)| \neq 1$ *and* $\alpha(G) > |V(G)|/2$, *then* $\xi(G) \geq 2$, *whenever* $|V(G)|$ *is odd, and* $\xi(G) \geq 3$, *if* $|V(G)|$ *is even.*

Remark 5 *The graph G in Figure 5 shows that the bounds in Theorem 2.16 are tight. We may consider either* $k = r-1$ *or* $k = r$, *and correspondingly,* $n+k-1$ *will be even or odd.*



Figure 5: $\alpha(G) = p + r,\ \xi(G) = r + 1$.

3 König-Egerváry graphs and bipartite graphs

In the sequel we deal with König-Egerváry graphs, for which we show that the converse of Theorem 2.14 is also true. It is known that $\lfloor n/2 \rfloor + 1 \leq \alpha(G) + \mu(G) \leq n$ holds for any graph G with n vertices.

Lemma 3.1 *If* G *is a König-Egerváry graph, then* $\alpha(G) \geq \mu(G)$.

Proof. Since $\mu(G) \leq |V(G)|/2$ holds for any graph G, and $|V(G)| = \alpha(G) + \mu(G)$ is true for G a König-Egerváry graph, then clearly follows that $\alpha(G) \geq \mu(G)$ is valid in our premise on G. ■

Corollary 3.2 *If* G *is a König-Egerváry graph, then the following statements are true:*

(i) $\alpha(G) \geq |V(G)|/2$;
(ii) $\alpha(G) = |V(G)|/2$ *if and only if* G *has a perfect matching;*
(iii) $\alpha(G) > |V(G)|/2$, *whenever* $isol(G) \neq \emptyset$.

Combining Proposition 2.8 and Corollary 3.2 we may conclude that:

Proposition 3.3 *If G is an α^+-stable König-Egerváry graph with $|isol\,(G)| \neq 1$, i.e., $isol\,(G) \neq \emptyset$, then $\alpha(G) = |V(G)|/2$.*

Remark 6 *There are non-α^+-stable König-Egerváry graphs G, without isolated vertices, such that $\alpha(G) = |V(G)|/2$. For instance, the graph G in Figure 6 has $\alpha(G) = 3 = |V(G)|/2$ and $\xi(G) = 2$, i.e., G is not α^+-stable. Notice that G has perfect matchings.*



Figure 6: A Koenig-Egervary graph G with $\alpha(G) = |V(G)|/2$ and $\xi(G) \geq 2$.

Proposition 3.4 *If G is a König-Egerváry graph, then the following statements are equivalent:*

(i) $\alpha(G) > |V(G)|/2$;
(ii) G has no perfect matching;
(iii) G is non-quasi-regularizable;
(iv) $\xi(G) = |core(G)| > |N(core(G))|$.

Proof. Theorem 2.14 ensures that $(i) \Rightarrow (iv)$; the step $(iv) \Rightarrow (iii)$ is clear by Theorem 2.1; the implication $(iii) \Rightarrow (ii)$ follows from Theorem 2.1; and $(ii) \Rightarrow (i)$ is true according to Corollary 3.2. ■

Remark 7 *In general, (ii) implies neither (i) nor (iii); e.g., the graph K_3. The graph in Figure 4 shows that (i) does not always follow from (iii) or from (iv).*

Remark 8 *If G is a König-Egerváry graph with $isol(G) \neq \emptyset$, then all the assertions in Proposition 3.4 are valid, since (i) follows from Corollary 3.2.*

Theorem 3.5 *If G is a König-Egerváry graph, then the following statements are equivalent:*

(i) $\alpha(G) = |V(G)|/2$;
(ii) G has a perfect matching;
(iii) G is quasi-regularizable;
(iv) $|core(G)| \leq |N(core(G))|$.

Proof. It follows from Proposition 3.4 and Lemma 3.1. ■

Remark 9 *There are quasi-regularizable graphs without perfect matching; e.g., the graph in Figure 7.*



Figure 7: G is quasi-regularizable, but has no perfect matching.

Remark 10 *The equivalence (ii)* $\Leftrightarrow$ *(iii) in Theorem 3.5 (and comparison with Theorem 2.1) shows that Hall's marriage Theorem is also true for König-Egerváry graphs. Moreover, according to (iv), it is sufficient to check Hall's condition only for one specific stable set, namely for* $core(G)$.

Remark 11 *The graph in Figure 8 shows that there exist non-quasi-regularizable graphs satisfying the inequality* $|core(G)| \leq |N(core(G))|$.



Figure 8: G is not quasi-regularizable, and $|core\,(G)| = 3 \leq |N(core(G))| = 4$.

Proposition 3.6 *If* $G = (A, B, E)$ *is a bipartite graph with* $|isol\,(G)| \neq 1$*, then either (i)* $\xi(G) \geq 2$*, or (ii)* $\xi(G) = 0$ *and* $A, B \in \Omega(G)$.

Proof. By Corollary 3.2, G satisfies $\alpha(G) \geq |V\,(G)|\,/2$. If $\alpha(G) > |V\,(G)|\,/2$, then $|isol\,(G)| \neq 1$ assures that $\xi(G) \geq 2$ holds, by Theorem 2.11. If $\alpha(G) = |V\,(G)|\,/2$, then Corollary 3.2(ii) implies that $|A| = |B| = |V\,(G)|\,/2$. Hence, $A, B \in \Omega(G)$ and evidently $\xi(G) = 0$. ■

Remark 12 *The above result is not true for all König-Egerváry graphs. Moreover, for any even positive integer k there exists a König-Egerváry graph G of order k with* $\xi(G) = 1$. *This observation is illustrated in Figure 9, where* $core(G) = \{c_2\}$.

If the order of a König-Egerváry graph G is odd, then $\alpha(G) > |V\,(G)|\,/2$ *and consequently* $\xi(G) \geq 2$. *However, for every odd positive integer k there exists a graph G of size k with* $\xi(G) = 1$. *For instance, the graph* $H = (V(G) \cup \{c_3\}, E(G) \cup \{a_1c_3, b_1c_3\})$ *is of odd order* $k = 2p + 3$ *and* $core(H) = \{c_2\}$.

Theorem 3.7 *If G is a bipartite graph with* $\xi(G) = 1$*, then* $|isol(G)| = 1$ *and* $|V(G)| \equiv 1$ *mod* 2.



Figure 9: A Koenig-Egervary graph G of order $k = 2p + 2$ with $\xi(G) = 1$.

Proof. Proposition 3.6 ensures that $|isol(G)| = 1$. Then, Corollary 3.2 implies that $\alpha(G) > |V(G)|/2$, and consequently, $|V(G)| \equiv 1 \ mod \ 2$ is now true, according to Corollary 2.10. ■

As a consequence of Proposition 3.6 we obtain the following characterization of α^+-stable bipartite graphs.

Corollary 3.8 *[8] For a bipartite graph G with $|isol(G)| \neq 1$, the following assertions are equivalent:*
(i) G is α^+-stable;
(ii) G has a perfect matching;
(iii) G possesses two maximum stable sets that partition its vertex set.

Proof. $(i) \Rightarrow (ii), (iii)$ By Proposition 3.3, we get that $\alpha(G) = |V(G)|/2$. Hence, Theorem 3.5 implies (ii), and (iii) follows from Proposition 3.6.

$(iii) \Rightarrow (i)$ It is evident, by Theorem 1.1.

$(ii) \Rightarrow (iii)$ If G has a perfect matching, then Theorem 3.5 assures that $\alpha(G) = |V(G)|/2$, and this leads (see the proof of Proposition 3.6) to $A, B \in \Omega(G)$. ■

This result generalizes the corresponding statement for trees, given in [4].

Clearly, Proposition 3.4 is true for bipartite graphs as well, since they are König-Egerváry graphs. Moreover, using Corollary 3.8, we obtain a stronger form of this assertion.

Proposition 3.9 *If G is a bipartite graph with $|isol(G)| \neq 1$, then*

$$\alpha(G) > |V(G)|/2 \text{ if and only if } \xi(G) \geq 2.$$

Proof. According to Theorem 2.11 (i), $|isol(G)| \neq 1$ and $\alpha(G) > |V(G)|/2$ implies that $\xi(G) \geq 2$ is true for general graphs.

Conversely, for G a bipartite graph with $|isol(G)| \neq 1$, we have to show that if $\xi(G) \geq 2$, then also $\alpha(G) > |V(G)|/2$. By Theorem 1.1, it follows that G is not α^+-stable, and then Corollary 3.8 implies that G has no perfect matching. Hence, by Corollary 3.2, we get that $\alpha(G) > |V(G)|/2$. ■

4 Conclusions

In this paper we have presented a number of relations connecting $\alpha(G)$ and $\xi(G)$, whenever $\alpha(G) > |V(G)|/2$, with emphasis on König-Egerváry graphs and bipartite graphs, for which some sufficient conditions are also necessary. It would be interesting to see which of these results can be transferred to graphs satisfying $\alpha(G) = |V(G)|/2$ or less. The special case of König-Egerváry graphs could offer a promising start, since for them the condition $\alpha(G) \leq |V(G)|/2$ is equivalent to $\alpha(G) = |V(G)|/2$. We also have shown that a necessary and sufficient condition for a König-Egerváry graph G to have a perfect matching reads as $|core(G)| \leq |N(core(G))|$. The following challenging problem seems to be of algorithmic nature: how to find $core(G)$, at least for König-Egerváry graphs.

Fugure 10 suggests the following questions: what must be added to $\xi(G) \geq 2$, in order to get $\alpha(G) > |V(G)|/2$ for, at least, König-Egerváry graphs; when the inequality $|core(G)| > |N(core(G))|$ can be obtained from $\xi(G) \geq 2$?



Figure 10: A scheme of interconnections between the main findings of the paper.

5 Acknowledgment

The authors thank Professor Uri Peled for his careful reading of this manuscript, and also for his suggestions and comments.

References

[1] C. Berge, *Some common properties for regularizable graphs, edge-critical graphs and B-graphs*, in "Graph Theory and Algorithms Lecture Notes in Computer Science", Vol. **108**, Springer-Verlag, Berlin, 1980, 108-123 .

[2] R. W. Deming, *Independence numbers of graphs - an extension of the König-Egerváry theorem*, Discrete Mathematics **27** (1979), 23-33.

[3] E. Egerváry, *On combinatorial properties of matrices*, Matematikai Lapok **38** (1931), 16-28.

[4] G. Gunther, B. Hartnell, and D. F. Rall, *Graphs whose vertex independence number is unaffected by single edge addition or deletion*, Discrete Applied Mathematics **46** (1993), 167-172.

[5] T. W. Haynes, L. M. Lawson, R. C. Brigham and R. D. Dutton, *Changing and unchanging of the graphical invariants: minimum and maximum degree, maximum clique size, node independence number and edge independence number*, Congresus Numerantium **72** (1990), 239-252.

[6] P. L. Hammer, P. Hansen, and B. Simeone, *Vertices belonging to all or to no maximum stable sets of a graph*, SIAM Journal of Algebraic Discrete Methods **3** (1982) 511-522.

[7] D. König, *Graphen und matrizen*, Matematikai Lapok **38** (1931), 116-119.

[8] V. E. Levit and E. Mandrescu, *On α-stable graphs*, Congressus Numerantium **124** (1997), 33-46.

[9] V. E. Levit and E. Mandrescu, *The structure of α-stable graphs*, The Third Krakow Conference On Graph Theory, Krakow University, Kazimierz Dolny, Poland (1997), math.CO/**9911227** (1999).

[10] V. E. Levit and E. Mandrescu, *On α^+-stable König-Egerváry graphs*, The Ninth SIAM Conference on Discrete Mathematics, University of Toronto, Canada (1998), math.CO/**9912022** (1999).

[11] V. E. Levit and E. Mandrescu, *Well-covered and König-Egerváry graphs*, Congressus Numerantium **130** (1998), 209-218.

[12] L. Lovasz and M. D. Plummer, "*Matching Theory*", Annals of Discrete Math. **29** (1986), North-Holland.

[13] F. Sterboul, *A characterization of the graphs in which the transversal number equals the matching number*, Journal of Combinatorial Theory Series B **27** (1979), 228-229.

[14] J. Zito, *The structure and maximum number of maximum independent sets in trees*, Journal of Graph Theory **15** (1991) 207-221.